

# Simplicity and complexity in the light-front vacuum

*Work done in collaboration with S.S. Chabysheva.*

*Supported in part by the Minnesota Supercomputing Institute and  
the IBEST Computational Resources Core at the University of Idaho.*

John R. Hiller

`jhiller@uidaho.edu`

Department of Physics

University of Idaho

# Introduction

- the light-front (LF) vacuum is famously considered trivial, while the equal-time (ET) vacuum is not
- there are several indications that this is not really the case
  - most recently, zero-mode contributions have been proposed for twist-3 GPDs  
[Aslan & Burkardt, PRD 101, 016010 (2020);  
X. Ji, NPB 960, 115181 (2020)]
- but what is missing?
- the key is that matrix elements of transitions to and from the Fock vacuum are not necessarily zero and terms usually dropped from LF Hamiltonians need to be kept
- this leads to vacuum bubbles and tadpoles, which we explore in a nonperturbative context


# Outline

- light-front quantization
- evidence for a nontrivial vacuum
- restoration of vacuum transitions
- vacuum bubbles for free & shifted scalars
- tadpoles and bubbles in  $\phi^4$  theory
- vacuum energy subtraction in quenched scalar Yukawa theory
- summary

# Light-front coordinates

Dirac, RMP 21, 392 (1949).

- time:  $x^+ = t + z$
- space:  $\underline{x} = (x^-, \vec{x}_\perp)$ ,  $x^- \equiv t - z$ ,  $\vec{x}_\perp = (x, y)$
- energy:  $p^- = E - p_z$
- momentum:  $\underline{p} = (p^+, \vec{p}_\perp)$ ,  $p^+ \equiv E + p_z$ ,  $\vec{p}_\perp = (p_x, p_y)$
- mass-shell condition:  $p^2 = m^2 \Rightarrow p^- = \frac{m^2 + p_\perp^2}{p^+}$



# Mass eigenvalue problem

- Pauli and Brodsky, PRD **32**, 1993 (1985); 2001 (1985)

$$\mathcal{P}^- |\underline{P}\rangle = \frac{M^2 + P_\perp^2}{P^+} |\underline{P}\rangle, \quad \mathcal{P} |\underline{P}\rangle = \underline{P} |\underline{P}\rangle.$$


- no spurious vacuum contributions to eigenstates
  - $p^+ = \sqrt{m^2 + p_\perp^2 + p_z^2} + p_z > 0$  for all massive particles
  - cannot produce particles from vacuum and still conserve  $p^+$
  - Fock vacuum  $|0\rangle$  is the physical vacuum
- boost-invariant separation of internal and external momenta
  - longitudinal momentum fractions  $x_i \equiv p_i^+ / P^+$
  - relative transverse momenta  $\vec{k}_{i\perp} \equiv \vec{p}_{i\perp} - x_i \vec{P}_\perp$

# $\phi^4$ critical couplings compared

Method	$g_c \equiv \lambda_c / (24\mu^2)$	Reported by
LF sym. polys.	$1.1 \pm 0.03$	Burkardt, SSC & JRH
DLCQ	1.38	Harindranath & Vary
Quasi-sparse eigenvector	2.5	Lee & Salwen
Density matrix	2.4954(4)	Sugihara
Lattice	$2.70 \begin{cases} +0.025 \\ -0.013 \end{cases}$	Schaich & Loinaz
	$2.79 \pm 0.02$	Bosetti <i>et al.</i>
Uniform matrix product	2.766(5)	Milsted <i>et al.</i>
Renorm. H trunc.	2.97(14)	Rychkov & Vitale

Systematic difference between LF (top) and ET (bottom).

# Troublesome tadpole



Contributes to ET self-energy, but not LF.

# Mass renormalization

Bare mass renormalized by tadpole contributions in ET quantization but not in LF quantization  
[M. Burkardt, PRD 47, 4628 (1993)]

$$\mu_{\text{LF}}^2 = \mu_{\text{ET}}^2 + \lambda \left[ \langle 0 | \frac{\phi^2}{2} | 0 \rangle - \langle 0 | \frac{\phi^2}{2} | 0 \rangle_{\text{free}} \right].$$

The vev's of  $\phi^2$  resum the tadpole contributions; the subscript *free* indicates the vev with  $\lambda = 0$ .

→ need to calculate vev's

[M. Burkardt, SSC, and JRH, PRD 94, 065006 (2016)]

$$\langle 0 | \frac{\phi^2}{2} | 0 \rangle \rightarrow \frac{1}{2} \langle 0 | \phi(\epsilon^+, \epsilon^-) \int_0^\infty dP \sum_n |\psi_n(P)\rangle \langle \psi_n(P) | \phi(0, 0) | 0 \rangle.$$

$$\phi(\epsilon^+, \epsilon^-) = e^{i\mathcal{P}^- \epsilon^+ / 2} \phi(0, \epsilon^-) e^{-i\mathcal{P}^- \epsilon^+ / 2}.$$



# ET to LF interpolation

K. Hornbostel, Phys. Rev. D **45**, 3781 (1992);

B. Ma and C.R. Ji, arXiv:2105.09388, refs and Thurs 14:40

- $x^\pm = \frac{1}{\sqrt{2}}[\sqrt{1 \pm ct} \pm \sqrt{1 \mp cz}] \rightarrow \text{ET: } c = 1, x^\pm = t, -z$

- LF:  $c = 0, x^\pm = (t \pm z)/\sqrt{2}$  as usual, modulo  $\sqrt{2}$

- $p_\pm = \frac{1}{\sqrt{2}}[\sqrt{1 \pm cE} \mp \sqrt{1 \mp cp_z}]$

- $p \cdot x = p_+ x^+ + p_- x^-$

- $\mu^2 = E^2 - p_z^2 = cp_+^2 - cp_-^2 + 2sp_+p_-, s \equiv \sqrt{1 - c^2}$

- positive root  $\rightarrow p_+ = [\sqrt{p_-^2 + c\mu^2 - sp_-}]/c$

- compute for arbitrary  $c$  and study limit

- SSC and JRH, Phys. Rev. D **102**, 116010 (2020)

- obtain consistency with equal-time


- LF appears inequivalent to  $c \rightarrow 0$  limit

# Nontrivial vacuum bubbles

- Collins [arXiv:1801.03960] considered the one-loop self-energy in  $\phi^3$  theory and the related invariant function

$$\Pi(p^2) = -\frac{1}{8\pi^2} \int \frac{d^2k}{[k^2 - \mu^2 + i\epsilon][(p-k)^2 - \mu^2 + i\epsilon]}.$$

- the one-loop bubble is obtained in the  $p^2 \rightarrow 0$  limit;  
 $\Pi(0) = -i/8\pi\mu^2$ .



- naive LF approach gives zero for the bubble.
- ET to LF transition agrees with ET.

# Proposed resolution

- restore vacuum transitions to LF quantization

- $$\mathcal{P}_{\text{free}}^- = \int dx^- : \frac{1}{2} \mu^2 \phi^2 := \int dp^+ \frac{\mu^2}{p^+} a^\dagger(p^+) a(p^+) \\ + \frac{\mu^2}{2} \int \frac{dp_1^+ dp_2^+}{\sqrt{p_1^+ p_2^+}} \delta(p_1^+ + p_2^+) [a(p_1^+) a(p_2^+) + a^\dagger(p_1^+) a^\dagger(p_2^+)]$$

- standard practice is to drop the last two terms, because  $p_i^+$  forced to zero
- instead keep; then have tadpoles and bubbles
  - matrix elements of such terms need not be zero
- need to regulate bubbles
  - perturbative calculations allow classification and exclusion of such contributions
  - nonperturbative calculations do not
  - $\delta \rightarrow \delta_\epsilon$  with  $\epsilon$  a width to be taken to zero
  - ‘ephemeral’ modes

# Free vacuum


- generalized coherent state:  $|\text{vac}\rangle_0 = \sqrt{Z} e^{\alpha A^\dagger} |0\rangle$ ,  
where  $A^\dagger \equiv \int \frac{dp_1 dp_2}{\sqrt{p_1 p_2}} \frac{f(p_1, p_2)}{\frac{1}{p_1} + \frac{1}{p_2}} a^\dagger(p_1) a^\dagger(p_2)$ .  
(have dropped plus superscript for brevity)
- solves  $\mathcal{P}_{\text{free}}^- |\text{vac}\rangle_0 = P_{\text{vac}}^- |\text{vac}\rangle_0$   
if  $\alpha f(p_1, p_2) = -\frac{1}{2} \delta_\epsilon(p_1 + p_2)$
- with  $P_{\text{vac}}^- = -\frac{\mu^2}{2} \int dQ \delta_\epsilon(Q)^2 = -\frac{\mu^2 L}{16\pi}$
- where  $\int dQ \delta_\epsilon(Q)^2 \rightarrow \delta(0) \int_0^\infty \delta(Q) dQ = \frac{1}{2} \frac{L}{4\pi}$   
and  $L = \int dx^- = 4\pi \delta(0)$  is the light-front volume
- a massive state  $a^\dagger(P) |\text{vac}\rangle_0$  is then an eigenstate of  $\mathcal{P}_{\text{free}}^- + \frac{\mu^2 L}{16\pi}$  with eigenvalue  $\frac{M^2}{P}$ 
  - with  $\epsilon \ll P$ , the ephemeral modes are disjoint

# Shifted scalar

- $\phi \rightarrow \phi + v$
- $\mathcal{L} = \mathcal{L}_0 - \mu^2 v \phi - \frac{1}{2} \mu^2 v^2$
- $\phi(x^+ = 0, x^-) = \int \frac{dp}{\sqrt{4\pi p}} \left\{ a(p) e^{-ipx^-/2} + a^\dagger(p) e^{ipx^-/2} \right\}$
- $\mathcal{P}_{\text{int}}^- = \int dx^- [\mu^2 v \phi + \frac{1}{2} \mu^2 v^2]$   
 $= \sqrt{4\pi} \mu^2 v \int \frac{dp}{\sqrt{p}} \delta_\epsilon(p) [a(p) + a^\dagger(p)] + \frac{1}{2} \mu^2 v^2 L$
- define  $B = -v \int dp \sqrt{4\pi p} \delta_\epsilon(p) a(p)$ ;  
then  $e^{B^\dagger} \mathcal{P}_{\text{free}}^- e^{-B^\dagger} = \mathcal{P}_{\text{free}}^- + \mathcal{P}_{\text{int}}^-$
- vacuum of the shifted Hamiltonian:  $|\text{vac}\rangle = e^{B^\dagger} |\text{vac}\rangle_0$
- vev:  $\langle \text{vac} | \phi(0) | \text{vac} \rangle = -v$

# Tadpoles and bubbles in $\phi^4$ theory

- light-front Hamiltonian density:  $\mathcal{H} = \frac{1}{2}\mu^2\phi^2 + \frac{\lambda}{4!}\phi^4$
  - light-front Hamiltonian:  $\mathcal{P}^- = \mathcal{P}_{\text{free}}^- + \mathcal{P}_{\text{eph}}^- + \dots$
- $$\mathcal{P}_{\text{eph}}^- = \frac{\lambda}{24} \int \frac{\prod_i^4 dp_i}{4\pi\sqrt{\prod_i^4 p_i}} \delta_\epsilon(\sum_i^4 p_i) [\prod_i^4 a(p_i) + \prod_i^4 a^\dagger(p_i)]$$
- focus on leading tadpole and bubble contributions



- Fock-state expansion:  $|\psi(P)\rangle = \psi_1 a^\dagger(P)|0\rangle + \dots + \int \prod_i^5 dp_i \delta(P - \sum_i^5 p_i) \psi_5(p_1, \dots, p_5) \frac{1}{5!} \prod_i^5 a^\dagger(p_i)|0\rangle + \dots$

# Equations for wave functions

- the eigenvalue problem:

$$(\mathcal{P}_{\text{free}}^- + \mathcal{P}_{\text{int}}^-)|\psi(P)\rangle = \left(\frac{M^2}{P} + P_{\text{vac}}^-\right)|\psi(P)\rangle$$

$$[P_{\text{vac}}^- \text{ obtained from solving } (\mathcal{P}_{\text{free}}^- + \mathcal{P}_{\text{int}}^-)|\text{vac}\rangle = P_{\text{vac}}^-|\text{vac}\rangle]$$

- projection onto Fock sectors

$$\frac{\mu^2}{P}\psi_1 + \frac{\lambda}{\sqrt{24}} \int \frac{\prod_i^4 dp_i}{4\pi\sqrt{\prod_i^4 p_i}} \delta_\epsilon(\sum_i^4 p_i) \psi_5(p_1, \dots, p_5) = \left(\frac{M^2}{P} + P_{\text{vac}}^-\right) \psi_1,$$

$$\left(\sum_i^5 \frac{\mu^2}{p_i}\right) \psi_5 + \frac{\lambda}{24} \frac{1}{5} \left[ \frac{\delta_\epsilon(\sum_i^4 p_i)}{4\pi\sqrt{\prod_i^4 p_i}} + (p_5 \leftrightarrow p_1, p_2, p_3, p_4) \right] \psi_1$$

$$+ 20 \frac{\lambda}{4} \int \frac{dp'_1 dp'_2}{4\pi\sqrt{p_1 p_2 p'_1 p'_2}} \delta(p_1 + p_2 - p'_1 - p'_2) \psi_5(p'_1, p'_2, p_3, p_4, p_5) = \frac{M^2}{P} \psi_5$$

- sector-dependent energy shift: no  $P_{\text{vac}}^-$  in the top sector

# Iterative solution

- second equation solved iteratively w.r.t. self-coupling of the five-constituent Fock state
  - this corresponds to a diagrammatic expansion
- substitute into first equation

- leading term generates the vacuum bubble

$$\int \frac{\prod_i^5 dp_i}{\prod_i^4 p_i} \delta(P - \sum_i^5 p_i) \frac{\delta_\epsilon(\sum_i^4 p_i)^2}{\frac{M^2}{P} - \sum_i^5 \frac{\mu^2}{p_i}}$$
$$\sim - \int \delta_\epsilon(Q)^2 \frac{dQ}{\mu^2} \int \frac{\prod_i^4 dx_i}{\prod_i^4 x_i} \delta(1 - \sum_i^4 x_i)$$

- diverges as  $\epsilon \rightarrow 0$ , proportional to  $\delta(0) = L/4\pi$
- canceled by bubble in  $P_{\text{vac}}^-$
- second term, where the self interaction acts once, produces the tadpole (next slide)



# Evaluation of tadpole

$$\int \frac{\prod_i^5 dp_i \delta_\epsilon(\sum_i^4 p_i) \frac{\delta(P - \sum_i^5 p_i)}{\frac{M^2}{P} - \sum_i^5 \frac{\mu^2}{p_i}}}{\sqrt{\prod_i^4 p_i}} \int \frac{dp'_1 dp'_2}{\sqrt{p_4 p_5 p'_1 p'_2}} \frac{\delta(p_4 + p_5 - p'_1 - p'_2)}{\frac{M^2}{P} - \sum_i^3 \frac{\mu^2}{p_i} - \frac{\mu^2}{p'_1} - \frac{\mu^2}{p'_2}} \frac{\delta_\epsilon(\sum_i^3 p_i + p'_1)}{\sqrt{\prod_i^3 p_i p'_1}}$$

- $\delta(P - \sum_i^5 p_i)$  reduces to  $\delta(P - p_5)$ 
  - used to do the  $p_5$  integral
- $\delta(p_4 + p_5 - p'_1 - p'_2)$  becomes  $\delta(p_4 + P - p'_1 - p'_2)$ 
  - used to do the  $p'_2$  integral
- $\delta_\epsilon(\sum_i^3 p_i + p'_1)$  can be written  $\delta_\epsilon(p_4 - p'_1)$ 
  - used to do the  $p'_1$  integral

- these leave 
$$\int \frac{\prod_i^4 dp_i}{\prod_i^4 p_i} \frac{1}{p_4 P} \frac{\delta_\epsilon(\sum_i^4 p_i)}{\left[\frac{M^2}{P} - \sum_i^4 \frac{\mu^2}{p_i} - \frac{\mu^2}{P}\right]^2}$$

$$\sim \frac{1}{P} \int_0^\infty \delta(Q) dQ \int \frac{\prod_i^4 dx_i}{(\prod_i^4 x_i) x_4 \left(\sum_i^4 \frac{\mu^2}{x_i}\right)^2}$$

# Tadpole self-energy correction

$$\sim \frac{1}{P} \int_0^\infty \delta(Q) dQ \int \frac{\prod_i^4 dx_i}{(\prod_i^4 x_i) x_4 \left( \sum_i^4 \frac{\mu^2}{x_i} \right)^2}$$

- finite and inversely proportional to  $P$ 
  - a light-front self-energy correction
- in a nonperturbative calculation, these contributions cannot be separated.
  - regulate bubbles via  $\delta \rightarrow \delta_\epsilon$
  - solve separate eigenproblems for the vacuum and the massive states
  - carry out the  $P_{\text{vac}}^-$  subtraction
  - take the width parameter  $\epsilon$  to zero.
- with interactions, ephemeral modes no longer disjoint
  - for strong coupling, wave functions are broad
  - may have edge effects (aka zero modes)

# Quenched scalar Yukawa theory

- simplest nonperturbative case (beyond shift)

- Lagrangian:

$$\mathcal{L} = |\partial_\mu \chi|^2 - m^2 |\chi|^2 + \frac{1}{2} (\partial_\mu \phi)^2 - \frac{1}{2} \mu^2 \phi^2 - g \phi |\chi|^2$$

- quenched light-front 2D Hamiltonian:

$$\mathcal{P}^- = \int dx^- \mathcal{H} = \mathcal{P}_{\text{free}}^- + \mathcal{P}_{\text{int}}^-$$

$$\begin{aligned} \mathcal{P}_{\text{free}}^- = & \int dp \frac{m^2}{p} \left[ c_+^\dagger(p) c_+(p) + c_-^\dagger(p) c_-(p) \right] + \int dq \frac{\mu^2}{q} a^\dagger(q) a(q) \\ & + \frac{\mu^2}{2} \int \frac{dq_1 dq_2}{\sqrt{q_1 q_2}} \delta_\epsilon(q_1 + q_2) \left[ a(q_1) a(q_2) + a^\dagger(q_1) a^\dagger(q_2) \right], \end{aligned}$$

$$\mathcal{P}_{\text{int}}^- =$$

$$g \int \frac{dp dq}{\sqrt{4\pi p q (p+q)}} \left\{ \left[ c_+^\dagger(p+q) c_+(p) + c_-^\dagger(p+q) c_-(p) \right] a(q) + \text{h.c.} \right\}$$

- quenching suppresses ephemeral modes for the complex scalar

- leaving only those of the neutral scalar

# Eigenproblem and Fock expansion

- charge-zero sector corresponds to the free scalar
  - provides the needed subtraction of  $P_{\text{vac}}^-$  for calculations in the charge-one sector
  - $P_{\text{vac}}^- = -\frac{\mu^2 L}{16\pi} = -\frac{\mu^2}{2} \int_0^\infty dQ \delta_\epsilon(Q)^2$
- charge-one sector:  $\mathcal{P}^- |\psi(P)\rangle = \left( \frac{M^2}{P} + P_{\text{vac}}^- \right) |\psi(P)\rangle$ 
  - eigenstate is a single complex scalar dressed by a cloud of neutrals
  - keeping only the first three Fock sectors

$$|\psi(P)\rangle = \psi_0 c_+^\dagger(P) |0\rangle + \int dq dp \delta(P - q - p) \psi_1(q) a^\dagger(q) c_+^\dagger(p) |0\rangle + \int dq_1 dq_2 dp \delta(P - q_1 - q_2 - p) \psi_2(q_1, q_2) \frac{1}{\sqrt{2}} a^\dagger(q_1) a^\dagger(q_2) c_+^\dagger(p) |0\rangle$$

# Coupled equations for wave functions

$$\begin{aligned} \frac{m^2}{P} \psi_0 + \frac{g}{\sqrt{4\pi}} \int_0^P \frac{dq \psi_1(q)}{\sqrt{qP(P-q)}} + \frac{\mu^2}{\sqrt{2}} \int \frac{dq_1 dq_2}{\sqrt{q_1 q_2}} \delta_\epsilon(q_1 + q_2) \psi_2(q_1, q_2) \\ = \left( \frac{M^2}{P} + P_{\text{vac}}^- \right) \psi_0, \end{aligned}$$

$$\begin{aligned} \left( \frac{\mu^2}{q} + \frac{m^2}{P-q} \right) \psi_1(q) + \frac{g}{\sqrt{4\pi}} \frac{\psi_0}{\sqrt{qP(P-q)}} \\ + \sqrt{2} \frac{g}{\sqrt{4\pi}} \int_0^{P-q} \frac{dq' \psi_2(q, q')}{\sqrt{q'(P-q)(P-q-q')}} = \frac{M^2}{P} \psi_1(q), \end{aligned}$$

and

$$\begin{aligned} \left( \frac{\mu^2}{q_1} + \frac{\mu^2}{q_2} + \frac{m^2}{P-q_1-q_2} \right) \psi_2(q_1, q_2) + \frac{\mu^2}{\sqrt{2}} \delta_\epsilon(q_1 + q_2) \frac{\psi_0}{\sqrt{q_1 q_2}} \\ + \frac{1}{\sqrt{2}} \frac{g}{\sqrt{4\pi}} \left[ \frac{\psi_1(q_1)}{\sqrt{q_2(P-q_1)(P-q_1-q_2)}} + \frac{\psi_1(q_2)}{\sqrt{q_1(P-q_2)(P-q_1-q_2)}} \right] \\ = \frac{M^2}{P} \psi_2(q_1, q_2) \end{aligned}$$

Again, the  $P_{\text{vac}}^-$  shift is taken to be sector dependent.

# Energy subtraction

- $\psi_2$  contains a piece proportional to  $\delta_\epsilon(q_1 + q_2)$

$$\begin{aligned}\psi_2 &\sim \left[ \frac{M^2}{P} - \frac{\mu^2}{q_1} - \frac{\mu^2}{q_2} - \frac{m^2}{P - q_1 - q_2} \right]^{-1} \frac{\mu^2}{\sqrt{2}} \delta_\epsilon(q_1 + q_2) \frac{\psi_0}{\sqrt{q_1 q_2}} \\ &\sim -\frac{1}{\sqrt{2}} \frac{\sqrt{q_1 q_2}}{q_1 + q_2} \delta_\epsilon(q_1 + q_2) \psi_0\end{aligned}$$

- substitution into the 3rd term of the 1st equation yields

$$\frac{\mu^2}{\sqrt{2}} \int \frac{dq_1 dq_2}{\sqrt{q_1 q_2}} \delta_\epsilon(q_1 + q_2) \psi_2(q_1, q_2) = -\frac{\mu^2}{2} \int \frac{dq_1 dq_2}{q_1 + q_2} \delta_\epsilon(q_1 + q_2)^2 \psi_0$$

- this is just  $P_{\text{vac}}^- \psi_0$ , which then cancels from both sides
  - this severe truncation is simple enough to allow this subtraction by hand
  - in general, this would not be the case
- next step is to solve the coupled equations
  - the  $\delta_\epsilon$  are best interpreted in integrals
  - matrix elements in a suitable basis will provide this

# Summary

- the LF vacuum is nontrivial in general
  - matrix elements of vacuum transition operators are not all zero
  - the vacuum must be computed and its energy subtracted
  - Fock-state wave functions have vacuum contributions
- vacuum transitions induce tadpole and bubble contributions
  - tadpoles provide the ‘missing link’ between ET and LF values of the  $\phi_2^4$  critical coupling
    - a calculation needs to be done to check
  - bubbles can be regulated and subtracted
- next steps: QSY,  $\phi_2^4$  theories